

March 9th

No 11

An
Inaugural Essay
ON
Hepatitis or Inflammation of the Liver
by
Cullen Lockett
of
Georgia.

admitted March 23d - 1821

[Faint, illegible handwriting, likely bleed-through from the reverse side of the page.]

acco
To
open
solid
long
O
or
acul
part
fig
in
get
lic
is
and
m
cal

Hepatitis, or Inflammation of the Liver.

This disease is located in class *Pneustia*, under *Phlegmasia*, according to Cullen.

Hepatitis is of two kinds, the one acute, which exhibits the essential character of genuine inflammation; the other chronic, which exhibits symptoms of less violence as to their inflammatory tendency.

Acute Hepatitis.

Symptoms. It commences in general with a shivering or rigor, which is succeeded by febrile heat with pain either acute or dull, in the right hypochondriac region, shooting more particularly in the direction of the back; and extending to the right shoulder-blade, and is accompanied with a cough, oppression in respiration and difficulty of lying, except on the right side; yet instances now and then occur where the patient is unable to lie on either side, pressure on the region of the liver, where there is usually some tension, induces considerable increase of pain and tendency to cough, and the cough attending this disease is more generally dry than moist. The alimentary canal becomes early implicated in this disease, as is indicated by the nausea;

side
fella
anom
are
pres
tong
occu
and
a mo
at the
leaves
of Ap
atten
the
The
for a
radio
decim
this a
on the

sickness, hiccup and frequently with a vomiting of apparently bilious matter; the bowels being for the most part irregular and generally slow in their action; the excrements discharged are of an ash or white colour, indicating the want of bile; the urine of a deep saffron colour, small in quantity, and is often tinged with bile; there are loss of appetite, great watchfulness and occasional delirium, extreme thirst, costiveness with a strong, hard, and frequent pulse, from ninety to one hundred and twenty in a minute, and sometimes intermitting; the skin is hot and dry at the same time, and the tongue covered with a white, and sometimes a yellowish crust, extending to the mouth and fauces; dysentery of spirits, and, when the disease has continued for some days it is attended generally, with a jaundice colour of the skin, by reason of the pressure of the inflamed liver upon the *vena bilialis*.

This disease is occasionally complicated with anomalous symptoms; for the liver has not unfrequently been found after death to be, scathed or otherwise injured, without any marked indication of disease during the life of the patient, excepting dyspepsia or simple indigestion. This disease has been known also, to spread itself with very great violence on the head, sometimes on the penis and lower extremities.

young
 all
 as
 are
 the
 face
 in
 that
 to
 soul
 they
 to
 an
 drag
 in
 that
 perha

Men from their depraved habits, are more subject to hepatitis than women; but children, perhaps, are the most frequent subjects of its attack in miasmatic countries.

It may be observed here, that in inflammation of the liver as well as in inflammation of every other part of the body, the symptoms are more or less violent, according to the degree of sensibility in the part affected. If the inflammation attacks the convex anterior surface of the liver, so that the peritonaeum becomes affected, the pain is much increased by pressure, and some degree of tumor may be observed. If that part of the organ be diseased, which is more immediately contiguous to the diaphragm it gives rise to painful and difficult respiration, acute shooting pains in the thorax. These symptoms from the resemblance they bear to those of the inflammatory affections of the chest, are liable to be confounded with them; but are to be readily distinguished by an attention to the history and progress of the complaint, and other diagnostics, which I will, presently, more particularly mention. In some cases of hepatitis, the stomach is so extremely irritable, that violent retching and vomiting occur; symptoms influenced, perhaps, by the inflammation being in the vicinity of that organ.

(C
 mon
 five
 occa
 year
 the
 min
 spen
 to co
 by be
 the p
 he m
 alon
 E
 for
 of in
 of the
 Rep
 affec
 the

CAUSES — Besides the most usual exciting causes of inflammation situated in other parts of the body, inflammation of the liver is occasioned by intense summer heats; hence it is of most frequent occurrence in tropical climates; by mental emotions as envy and jealousy, by blows on the head, by violent exercise, especially in the sun, by long continued, intermittent and remittent fevers, by miasmata, by high living, by the intemperate use of vinous and spirituous liquors, more particularly the latter, by the sudden exposure to cold when the body has been sometime in a state of perspiration, by biliary concretions which sometimes form in the gall ducts preventing the free passage of the bile to its destination, by the suppression of an hæmorrhoidal discharge and by some of the derangements of the alimentary canal.

Diagnosis. — The diseases which are liable to be mistaken for hepatitis, are the inflammatory affections of the chest, inflammation of the stomach, spasm of the gall ducts, and the rheumatic affections of the muscles in the neighbouring parts.

Hepatitis is more liable to be confounded with the inflammatory affections of the chest, when it is situated in the membrane covering the upper convex surface of the liver, or the ligaments that connect it

to the
system
in so
on the
board
of the
a trial
mean
leading
to the
man
on the
of the
person
after
things
of the
by the
one of

to the diaphragm, or even in the upper part of its sarenchymatous texture; by reason of the enlargement, which the inflammation may induce in it, so as to extend in such a direction, as to occasion it to make pressure on the diaphragm, sufficient to diminish the cavity of the chest, so that there is a considerable restriction and pain kept up by the motion of the diaphragm in the act of respiration. Under such circumstances a troublesome cough comes on, so that the disease assumes in some measure the character of thoracic inflammation. But we may readily distinguish them, by the pain, in the former extending to the shoulder, the pallor of the countenance, the cough being unaccompanied by expectoration, and there being less dyspnoea. In hepatitis inflammation of the liver is distinguished from that of the stomach, by its not being accompanied with that extreme sense of heat and pain, with which that organ is affected, especially after taking anything into it, and without the immediate rejection of them; nor is the debility of the system so great in the inflammation of the liver as in that of the stomach.

Hepatitis may be distinguished from spasm of the gall ducts, by there being no nausea, the pain being permanent the pulse being one hundred and upwards in a minute, and the patient at least

pit
er
is
S
for
to the
in
of the
in
her
be
gro
for
in
sp
pre
flow
dix

preferring to keep the body in a strait and quiescent posture, where—as the greatest ease, when there is spasm, in the gall ducts is obtained by bending the body forwards on the knees.

And in the case of muscular pain there is little or no fever, the pain more diffused, frequently removing from one part of the abdomen to the other, and becomes more diffused by varying the posture of the body, it generally alternates with rheumatic pain in one or more joints of the body.

Terminations. Cholitis like inflammation situated in other parts of the body, may terminate by resolution, suppuration, coarctation, or gangrene, but it most frequently terminates (from being badly managed) in chronic hepatitis.

The tendency of hepatitis to resolution is known by the general mildness of the symptoms and their yielding to the proper remedies, particularly by there being little cough, dyspnea, vomiting, oppression or debility. When this disease terminates spontaneously and favourably, there is generally some preternatural evacuation that takes place, as hæmorrhagy, from the nose, or hæmorrhoids, particularly; sometimes by diarrhoea, erysipelas inflammation, perspiration, or a copious

(68)
see
see
of the

love
of the
see
on
four
deso-
pat.
citic
poet
and
these
in c
the
we

discharge of high coloured urine, depositing a red crystalline sediment. In some cases there has been such a premature increase of the bronchial secretions, accompanying the resolution of this disease, as to occasion suffocations.

The symptoms indicating the formation of matter, in the liver, are not always very obvious, however the most striking of them are a diminution of pain, a sense of pulsation, and weight in the right hypochondrium, especially when lying on the left side, frequent returning rigors and an accession of fever. When hepatitis terminates in suppuration, it may discharge itself in several ways, by the stomach, or some other part of the alimentary canal, into the cavity of the abdomen, the cavity of the chest or the substance of the lung itself, the bladder, the pericardium; through the parietes of the abdomen, and the matter sometimes passes through the bladder into the duodenum; these two last modes are, perhaps the most favourable manners, in which the matter can be discharged.

When hepatitis does not altogether yield to the treatment advised, but being in some measure subdued by it, we may then expect it to terminate in abscess, which is known



By the enlarged, indurated, liver, that is found upon examinations, and pointed out by a dull pain situated in the right hypochondriac region, extending to the shoulder; also by an uneasy sensation, in respiration, occasioned by the pressure of the diseased liver upon the diaphragm, and also by digestion in general not going on regular, from the incapacity of the diseased liver to secrete a sufficient quantity of bile.

It is sometimes the case, where the pain and inflammation have subsided very suddenly, and then succeeded by a low fluttering pulse, cold extremities, delirium, and death, so that there have been just reasons to suspect that this organ, may on some occasions, though much less frequently than others, become gangrenous.

Diseased Appearances of the Liver.

On the dissection of those who die of hepatitis, it will be found that the liver has formed adhesions to the neighbouring parts, or that the membranes and liver, are more or less inflamed or altered in their appearances; in a state of suppuration, congestion, scirrhus, or as has sometimes been found gangrenous.

804
 805
 806
 807
 808
 809
 810
 811
 812
 813
 814
 815
 816
 817
 818
 819
 820
 821
 822
 823
 824
 825
 826
 827
 828
 829
 830
 831
 832
 833
 834
 835
 836
 837
 838
 839
 840
 841
 842
 843
 844
 845
 846
 847
 848
 849
 850
 851
 852
 853
 854
 855
 856
 857
 858
 859
 860
 861
 862
 863
 864
 865
 866
 867
 868
 869
 870
 871
 872
 873
 874
 875
 876
 877
 878
 879
 880
 881
 882
 883
 884
 885
 886
 887
 888
 889
 890
 891
 892
 893
 894
 895
 896
 897
 898
 899
 900
 901
 902
 903
 904
 905
 906
 907
 908
 909
 910
 911
 912
 913
 914
 915
 916
 917
 918
 919
 920
 921
 922
 923
 924
 925
 926
 927
 928
 929
 930
 931
 932
 933
 934
 935
 936
 937
 938
 939
 940
 941
 942
 943
 944
 945
 946
 947
 948
 949
 950
 951
 952
 953
 954
 955
 956
 957
 958
 959
 960
 961
 962
 963
 964
 965
 966
 967
 968
 969
 970
 971
 972
 973
 974
 975
 976
 977
 978
 979
 980
 981
 982
 983
 984
 985
 986
 987
 988
 989
 990
 991
 992
 993
 994
 995
 996
 997
 998
 999
 1000
 1001
 1002
 1003
 1004
 1005
 1006
 1007
 1008
 1009
 1010
 1011
 1012
 1013
 1014
 1015
 1016
 1017
 1018
 1019
 1020
 1021
 1022
 1023
 1024
 1025
 1026
 1027
 1028
 1029
 1030
 1031
 1032
 1033
 1034
 1035
 1036
 1037
 1038
 1039
 1040
 1041
 1042
 1043
 1044
 1045
 1046
 1047
 1048
 1049
 1050
 1051
 1052
 1053
 1054
 1055
 1056
 1057
 1058
 1059
 1060
 1061
 1062
 1063
 1064
 1065
 1066
 1067
 1068
 1069
 1070
 1071
 1072
 1073
 1074
 1075
 1076
 1077
 1078
 1079
 1080
 1081
 1082
 1083
 1084
 1085
 1086
 1087
 1088
 1089
 1090
 1091
 1092
 1093
 1094
 1095
 1096
 1097
 1098
 1099
 1100
 1101
 1102
 1103
 1104
 1105
 1106
 1107
 1108
 1109
 1110
 1111
 1112
 1113
 1114
 1115
 1116
 1117
 1118
 1119
 1120
 1121
 1122
 1123
 1124
 1125
 1126
 1127
 1128
 1129
 1130
 1131
 1132
 1133
 1134
 1135
 1136
 1137
 1138
 1139
 1140
 1141
 1142
 1143
 1144
 1145
 1146
 1147
 1148
 1149
 1150
 1151
 1152
 1153
 1154
 1155
 1156
 1157
 1158
 1159
 1160
 1161
 1162
 1163
 1164
 1165
 1166
 1167
 1168
 1169
 1170
 1171
 1172
 1173
 1174
 1175
 1176
 1177
 1178
 1179
 1180
 1181
 1182
 1183
 1184
 1185
 1186
 1187
 1188
 1189
 1190
 1191
 1192
 1193
 1194
 1195
 1196
 1197
 1198
 1199
 1200
 1201
 1202
 1203
 1204
 1205
 1206
 1207
 1208
 1209
 1210
 1211
 1212
 1213
 1214
 1215
 1216
 1217
 1218
 1219
 1220
 1221
 1222
 1223
 1224
 1225
 1226
 1227
 1228
 1229
 1230
 1231
 1232
 1233
 1234
 1235
 1236
 1237
 1238
 1239
 1240
 1241
 1242
 1243
 1244
 1245
 1246
 1247
 1248
 1249
 1250
 1251
 1252
 1253
 1254
 1255
 1256
 1257
 1258
 1259
 1260
 1261
 1262
 1263
 1264
 1265
 1266
 1267
 1268
 1269
 1270
 1271
 1272
 1273
 1274
 1275
 1276
 1277
 1278
 1279
 128

Treatment. As the symptoms of this disease vary as regards the climate in which it occurs, so the treatment must accordingly vary.

In the treatment of this disease, as it usually presents itself, we commence with copious venesection, from twenty to thirty ounces of blood, is to be taken at once, and the repetition of the operation, within twenty four hours, must be determined on by the degree of pain and fever with which the patient is affected, at the same time applying cups or leeches to the region of the liver, immediately to be followed by the application of a large blister over the part affected. In the meantime however, we should accompany the bloodletting by purging freely, with calomel, to be followed within eight or ten hours by a dose of Epsom salts, magnesia or some other laxative, or an emollient injection, which contribute to mitigate tormina and tenesmus. After the alimentary canal has been once freely evacuated, it should be kept in a soluble state by the following prescriptions;

R Calomel grs 6

Tartar Emetic grs 4


Quina grs ʒss M. f. t. Colur, is to be taken twice, in, four &c;

10. 10.
11. 11.
12. 12.
13. 13.
14. 14.
15. 15.
16. 16.
17. 17.
18. 18.
19. 19.
20. 20.
21. 21.
22. 22.
23. 23.
24. 24.
25. 25.
26. 26.
27. 27.
28. 28.
29. 29.
30. 30.
31. 31.
32. 32.
33. 33.
34. 34.
35. 35.
36. 36.
37. 37.
38. 38.
39. 39.
40. 40.
41. 41.
42. 42.
43. 43.
44. 44.
45. 45.
46. 46.
47. 47.
48. 48.
49. 49.
50. 50.
51. 51.
52. 52.
53. 53.
54. 54.
55. 55.
56. 56.
57. 57.
58. 58.
59. 59.
60. 60.
61. 61.
62. 62.
63. 63.
64. 64.
65. 65.
66. 66.
67. 67.
68. 68.
69. 69.
70. 70.
71. 71.
72. 72.
73. 73.
74. 74.
75. 75.
76. 76.
77. 77.
78. 78.
79. 79.
80. 80.
81. 81.
82. 82.
83. 83.
84. 84.
85. 85.
86. 86.
87. 87.
88. 88.
89. 89.
90. 90.
91. 91.
92. 92.
93. 93.
94. 94.
95. 95.
96. 96.
97. 97.
98. 98.
99. 99.
100. 100.

after which a laxative should be administered. The
laxative emetic and opium are given for the purpose of relieving the
hot and dry skin, which is generally attendant on this disease,
and to guard the bowels from irritation.

These remedies connected with a rigid enforcement of the antiphlogistic
regimen, constitute our main reliance in the commencement of the
phlegmon; and if the inflammation should readily yield to these,
nothing more directly for its management will be required, but
for the purpose of more resolutely eradicating the disease; it is
advisable for the patient to undergo a gentle mercurial punction,
which is to be continued for sometime with a mild regimenter diet.

But when the disease shall have continued for four or five days
without being at all subdued in its violence by the precursory
remedies, we may then expect suppuration of the liver shortly to
take place, but we must prevent it by producing a ptyalism as quick
as possible, and in order to fulfil our intention we must make use
of mercury to its greatest extent, both internally and externally; for
unfortunately at this critical period there seems to exist a morbid
such an susceptibility in the system to a mercurial ptyalism at the
time suppuration is about to take place or has taken place, that we are



by the administration of the best discriminating doses internally and by the most judicious incisions externally, occasionally aided in reducing the desired mercurial impressions.

Here the ipecac bath by determining it the surface will sometimes so far restore the balance of excitability and circulation, as to promote the action of mercury both from the internal and external surfaces of the body. But great care is to be taken to avoid a subsequent chill and a consequent recoil of the circulation which will be sure to aggravate all the symptoms, instead of relieving them. The action of mercury on the system is accelerated in some measure by causing the patient to swallow every night at bed time, a considerable quantity of warm diluting drinks, as thin water gruel &c.

The mercury may also be assisted in producing its effects by applying the nitro-muriatic acid bath of Mr. Seether the administration of the nitro-muriatic acid intervention in the quantity of one or two drachms a day to be taken in divided doses.

When all our endeavours to subdue the inflammation fails, and an abscess is about forming in the liver, which is marked by the pulse becoming softer, frequent shiverings, a lame nty pain, and a sensation of weight about the region of the liver, we must

a de
 and a
 his p
 for
 as
 strong
 he
 la
 a
 a
 so
 as to
 the
 as
 he
 of
 ge
 he
 can
 the

endeavour as much as possible, to promote the suppuration and its discharge through the parietics of the abdomen; for this purpose the usual application of mollient poultices and fomentations, must be immediately made, and duly persisted in, until the contents the abscess can be discharged by an incision through the external teguments by some skilful operator.

The opening should be made in the most depending part of the tumor, where a fluctuation is sometimes perceived. Having made a scalp cut through the external teguments, and reached the abscess, it may be pierced with a Trepan, and the matter gradually evacuated. If the liver has formed such adhesions to the peritoneum, as to prevent the pus from falling into the cavity of the abdomen the life of the patient may be preserved. The Peruvian bark in doses of half a drachm in powder, should be given four or five times a day during the formation of the abscess, and after it is opened, in order to support the strength of the patient, and to guard against the effects of purulent matter. The patient is to be supported with light nourishing food, and cooling, delicious drinks, carefully avoiding all heating substances, as in other inflammations.

The matter of abscess of the liver, may discharge itself in the



various manners before mentioned; which pass according to the adhesions that the liver forms, and the particular part in which the suppuration takes place; these cases are to be treated in the same general manner as the preceding ones; but they are much more apt to terminate fatally.

This, as the treatment of acute hepatitis as it occurs in the more temperate climates; but it will not answer in the tropical climates for it there runs into suppuration much quicker from the more violent action of the exciting causes, therefore the treatment consist in life depletion by the lancet, but a more active and early use of mercury for the purpose of completely putting the system under its influence.



Chronic Hepatitis

This form of the disease, as has been before observed, is the most frequent manner, in which acute hepatitis terminales have being badly managed; but it is occasionally induced by many of the same causes that excite the acute kinds, but they are milder in their effects, and it may thwart as the consequence of that disease; it is attended by most of the symptoms that characterize acute hepatitis, however they are less violent in their action, being accompanied with more sharpness and swelling, which are the most prominent diagnostics of the intensity of the disease.

The liver may exist in several states while exhibiting the symptoms of chronic hepatitis; as in a moderate state of inflammation in the membrane or substance of the liver, perhaps this is the most frequent manner in which it does exist while exhibiting these symptoms, could converted into cartilage; common induritis; very soft in its substance; very hard in its substance; worms in the liver; and the liver may be almost destroyed of every thing except its bloodvessels. We have no particular diagnostic by which we are enabled to distinguish these different states of the liver when thus united, therefore we are unable to point out a treatment for each one, but in the generality of cases the following will succeed best.

Be
1. 18
2. 18
3. 18
4. 18
5. 18
6. 18
7. 18
8. 18
9. 18
10. 18
11. 18
12. 18
13. 18
14. 18
15. 18
16. 18
17. 18
18. 18
19. 18
20. 18
21. 18
22. 18
23. 18
24. 18
25. 18
26. 18
27. 18
28. 18
29. 18
30. 18
31. 18
32. 18
33. 18
34. 18
35. 18
36. 18
37. 18
38. 18
39. 18
40. 18
41. 18
42. 18
43. 18
44. 18
45. 18
46. 18
47. 18
48. 18
49. 18
50. 18
51. 18
52. 18
53. 18
54. 18
55. 18
56. 18
57. 18
58. 18
59. 18
60. 18
61. 18
62. 18
63. 18
64. 18
65. 18
66. 18
67. 18
68. 18
69. 18
70. 18
71. 18
72. 18
73. 18
74. 18
75. 18
76. 18
77. 18
78. 18
79. 18
80. 18
81. 18
82. 18
83. 18
84. 18
85. 18
86. 18
87. 18
88. 18
89. 18
90. 18
91. 18
92. 18
93. 18
94. 18
95. 18
96. 18
97. 18
98. 18
99. 18
100. 18

Treatment. - It is sometimes required in the treatment of this disease, that we should make use of purgation, but in a very moderate degree and what is taken should be taken in a very moderate ~~large~~ quantity at a time. He should purge from the commencement of this disease, very copiously with calomel, to be given every night or two, to be followed next morning by a dose of epsom salt or magnesia.

Among the remedies in this complaint, mercury is the one on which we are principally to rely, given slowly and in small doses, so as to keep up a brassy taste in the mouth for some time, for the purpose of effectually promoting the secretion of bile, and excluding the excrementitious on the surface of the body; to increase the latter effect however, it has been found, useful to combine with it a small proportion of opium and tartar emetic, (as is recommended in the treatment of acute hepatitis) at the same time to guard the bowels from irritation.

The next most salutary process that is to be attended to, is to keep up a regular peristaltic motion of the bowels, and excite the mouths of the excretory ducts of the liver, which will tend to eliminate the rigid and depraved secretion from that organ itself.



There seems no medicine better calculated for that purpose than the following prescription:

R. Compound Extract Colocynthis.	ʒj
Calomel,	ʒj
Tartar. Emelic,	grs. iij
Oil Cloves,	ʒij ss
Common Syrup,	q. s. M.

Two of the above pills, taken occasionally at bed time, will move the bowels gently next morning, carry off disease, and promote healthy secretion of bile; and will be found to operate in a wonderful manner that mental dependency and long train of nervous symptoms so constantly attendant on this complaint.

Our attention at the same time is to be directed to the cutaneous discharges. This is never to be forced by heating or stimulating, but an immense habitus promoves, by the most gentle means, such as moderate exercise, particularly by gestation; by a sea voyage; swimming; with flannel next the skin, particularly in changeable climates. The assiduous and daily application of the flesh brush over the hypochondriac region, will be found to excite the healthy action of the biliary organs in no mean degree.



The tepid bath, using the utmost caution, in avoiding a subsequent chill, will evidently be serviceable. Blisters or the more permanent drain of a seton, in the side, where there is much local pain or uneasiness, may be had recourse to with advantage.

Another remedy which has acquired great reputation in the treatment of chronic hepatitis, is the nitric acid. It may be given in divided doses to the amount of from one to three drachms, in the course of twenty four hours. A pleasant formula is, to make it into a julep with gum Arabic and loaf sugar.

Scirrhus affections of the liver may be treated, in the same manner as chronic hepatitis.

The diet best adapted for persons labouring under hepatitis, is such as is nutritive and easy of digestion; and may be gradually improved until health is perfectly restored.

The who labours under diseased liver, and hopes to protract his existence, with any kind of comfort to himself, must abandon what are called the pleasures of the table, but which are in reality the bane of human health, and observe a rigid temperance with respect to diet.

of Se
Lett
His
and
for a
corri
of the

With respect to drinks, simple water is best; no
spirited liquors whatever should be drank.

Late hours and night air, are to be studiously avoided.

It is advisable to those who labour under affections of the liver,
and reside in tropical climates, for them to change their residence,
for a more temperate one; for, if they remain there, they are
continually exposed to one of the most frequent exciting causes
of those diseases.

